

REDUCED CLIQUE GRAPHS: A CORRECTION TO “CHORDAL GRAPHS AND THEIR CLIQUE GRAPHS”

DILLON MAYHEW AND ANDREW PROBERT

ABSTRACT. Galinier, Habib, and Paul introduced the reduced clique graph of a chordal graph G . The nodes of the reduced clique graph are the maximal cliques of G , and two nodes are joined by an edge if and only if they form a non-disjoint separating pair of cliques in G . In this case the weight of the edge is the size of the intersection of the two cliques. A clique tree of G is a tree with the maximal cliques of G as its nodes, where for any $v \in V(G)$, the subgraph induced by the nodes containing v is connected. Galinier et al. prove that a spanning tree of the reduced clique graph is a clique tree if and only if it has maximum weight, but their proof contains an error. We explain and correct this error.

In addition, we initiate a study of the structure of reduced clique graphs by proving that they cannot contain any induced cycle of length five (although they may contain induced cycles of length three or any even integer greater than two). We show that no cycle of length four or more is isomorphic to a reduced clique graph. We prove that the class of clique graphs of chordal graphs is not comparable to the class of reduced clique graphs of chordal graphs by providing examples that are in each of these classes without being in the other.

1. INTRODUCTION

We consider only simple graphs. A *chord* of a cycle is an edge that joins two vertices of the cycle without being in the cycle itself. A graph is *chordal* if any cycle with at least four vertices has a chord. A *clique* is a set of pairwise adjacent vertices. If S is a set of vertices and P is a path, then P is *S -avoiding* if no internal vertex of P is in S . Assuming that a and b are distinct vertices, an *ab -separator* is a set S of vertices not containing either a or b such that there is no S -avoiding path from a to b . If, in addition, S does not properly contain an ab -separator then it is a *minimal ab -separator*.

If G is a chordal graph, then $C(G)$ is the corresponding *clique graph* (also known as the *clique intersection graph*). The vertices of $C(G)$ are the maximal cliques of G , and two maximal cliques are adjacent in $C(G)$ if and only if they have a non-empty intersection. The vertices of the *reduced clique graph*, $C_R(G)$, are again the maximal cliques of G , but C and C' are adjacent in $C_R(G)$ if and only if $C \cap C' \neq \emptyset$ and C and C' form a *separating pair*: that is, there is no $(C \cap C')$ -avoiding path from a vertex in $C - C'$

to a vertex in $C' - C$. Note that the vertices of $C_R(G)$ are identical to the vertices of $C(G)$, and every edge of $C_R(G)$ is an edge of $C(G)$.




FIGURE 1. A chordal graph, its clique graph, and its reduced clique graph.

The reduced clique graph was introduced in [3] (where it is called a clique graph) and studied further in [5–8].

Let G be a graph, and let T be a tree whose vertices are the maximal cliques of G . If, for every $v \in V(G)$, the maximal cliques of G that contain v induce a connected subgraph of T , then T is a *clique tree*. Clique trees were introduced by Gavril [4], who proved that a graph has a clique tree exactly when it is chordal.

We weight each edge of $C_R(G)$ as follows: the edge joining cliques C and C' is weighted with $|C \cap C'|$. The following result is [3, Theorem 6].

Theorem 1.1. *Let G be a connected chordal graph. Let T be a spanning tree of $C_R(G)$. Then T is a clique tree if and only if it is a maximum-weight spanning tree.*

Although the statement of Theorem 1.1 is correct, the proof in [3, Theorem 6] is not. The issue arises in the proof that a maximum-weight spanning tree must be a clique tree. We illustrate the error by using the same argument to prove a false statement.

Non-theorem 1.2. *Let G be a chordal graph. Let C_0, C_1, \dots, C_n be the sequence of maximal cliques in a path of $C_R(G)$ where $n > 1$. Assume that there is a vertex v of G such that v is in $C_0 \cap C_n$, but in none of the cliques C_1, \dots, C_{n-1} . Then C_0 and C_n are adjacent in $C_R(G)$.*

Non-proof. Consider the subgraph G' of G induced by $C_0 \cup C_1 \cup \dots \cup C_n$. Thus G' is chordal. From [10, Corollary 2] we see that either v is a *simplicial vertex* (meaning that the neighbours of v in G' form a clique), or there is a pair, a, b , of vertices such that v belongs to a minimal ab -separator of G' . In the former case v is in a unique maximal clique of G' ([1, Theorem 3.1]). But C_0 and C_n are distinct maximal cliques of G' that contain v . Therefore

we can let S be a minimal ab -separator of G' , where v is in S . The proof of [2, Lemma 2.3] shows that there are two distinct maximal cliques, D_a and D_b , of G' such that D_a and D_b properly contain S , and $D_a - S$ is in the same connected component of $G' - S$ as a , while $D_b - S$ is in the same component as b . Thus D_a and D_b are maximal cliques of G' that contain v . But the only maximal cliques of G' that contain v are C_0 and C_n . Therefore we can assume without loss of generality that $D_a = C_0$ and $D_b = C_n$. Any path from a vertex of $C_0 - C_n$ to a vertex of $C_n - C_0$ must contain a vertex in $S = D_a \cap D_b = C_0 \cap C_n$. Therefore C_0 and C_n form a non-disjoint separating pair, so C_0 and C_n are adjacent in $C_R(G)$, as claimed. \square

We can see that this non-theorem is, indeed, not a theorem by examining Figure 1. Set C_0 , C_1 , and C_2 to be the maximal cliques $\{2, 3, 4, 6, 8, 9\}$, $\{1, 2, 3\}$, and $\{2, 3, 5, 7, 8, 9\}$, respectively. Thus C_0, C_1, C_2 is the vertex sequence of a path in $C_R(G)$. The vertex 8 is in $C_0 \cap C_2$, but not in C_1 . However C_0 and C_2 are not adjacent in $C_R(G)$. The error in the proof lies in the claim that “the only maximal cliques of G' that contain v are C_0 and C_n ”. This need not be true. Indeed, $\{2, 3, 4, 5, 8, 9\}$ is a maximal clique in the subgraph induced by $C_0 \cup C_1 \cup C_2$, and it contains 8, but it is not equal to either C_0 or C_2 . Exactly the same error appears in the proof of [3, Theorem 6]. Nonetheless, Theorem 1.1 is true, and we prove it in the next section.

2. REDUCED CLIQUE GRAPHS AND CLIQUE TREES

In [9] we will apply our main theorem to some matroid problems. For these purposes we would like to extend its scope somewhat. Instead of weighting the edges of $C_R(G)$ with sizes of intersections, we consider more general weightings.

Definition 2.1. Let G be a chordal graph. We consider a function σ which takes

$$\{\emptyset\} \cup \{C \cap C' : C, C' \text{ are distinct maximal cliques of } G\}$$

to non-negative integers. We insist that $\sigma(\emptyset) = 0$ and if X and X' are in the domain of σ and $X \subset X'$, then $\sigma(X) < \sigma(X')$. In such a case the function σ is a *legitimate weighting* of G . We weight an edge between C and C' with $\sigma(C \cap C')$.

Theorem 2.2. *Let G be a connected chordal graph and let σ be a legitimate weighting of G . Every clique tree is a spanning tree of $C_R(G)$ and every edge of $C_R(G)$ is contained in a clique tree. Moreover, a spanning tree of $C_R(G)$ is a clique tree if and only if it has maximum weight amongst all spanning trees.*

Note that the function that takes each intersection $C \cap C'$ to $|C \cap C'|$ is a legitimate weighting, so Theorem 2.2 does indeed imply Theorem 1.1. We now start proving the intermediate results required for the proof of Theorem 2.2.

Proposition 2.3. *Let G be a chordal graph, and let C and C' be maximal cliques of G . Let S be a set of vertices that contains $C \cap C'$. Let v_0, v_1, \dots, v_k be the vertex sequence of P , a shortest-possible S -avoiding path from a vertex in $C - C'$ to a vertex in $C' - C$. Then $(C \cap C') \cup \{v_i, v_{i+1}\}$ is a clique for each $i = 0, 1, \dots, k - 1$.*

Proof. If $C \cap C' = \emptyset$ then the result holds trivially, so we assume $C \cap C'$ is non-empty. Note that every vertex in $C \cap C'$ is adjacent to v_0 , and also to v_k , since these vertices are in $C - C'$ and $C' - C$. Now the result can only fail if there is a vertex $x \in C \cap C'$ that is not adjacent to v_i for some $i \in \{1, \dots, k - 1\}$. Let p be the largest integer such that $p < i$ and x is adjacent to v_p . Similarly, let q be the smallest integer such that $q > i$ and x is adjacent to v_q . Consider the cycle obtained by adding the edges $v_p x$ and $v_q x$ to v_p, v_{p+1}, \dots, v_q . This cycle contains the distinct vertices v_p, v_i, v_q , and x , so it must contain a chord. No chord can join two vertices in the path P , since P is as short as possible. Thus any chord is incident with x . But x is not adjacent to any of the vertices in v_{p+1}, \dots, v_{q-1} by the choice of p and q , so we have a contradiction. \square

Proposition 2.4. *Let G be a chordal graph, and let C and C' be maximal cliques of G where $C \cap C' \neq \emptyset$. If C and C' are not adjacent in $C_R(G)$, then they are joined by a path of $C_R(G)$ with vertex sequence C_0, C_1, \dots, C_s , where each $C_i \cap C_{i+1}$ properly contains $C \cap C'$.*

Proof. Assume this fails for C and C' , and they have been chosen so that $C \cap C'$ is as large as possible. Let S be $C \cap C'$. Because C and C' are not adjacent in $C_R(G)$, but $S \neq \emptyset$, it follows that there is an S -avoiding path from a vertex in $C - C'$ to a vertex in $C' - C$. Let v_0, v_1, \dots, v_k be the vertex sequence of such a path, where k is as small as possible. We assume v_0 is in $C - C'$ while v_k is in $C' - C$. We apply Proposition 2.3 and for each $i = 1, \dots, k$, we let D_i be a maximal clique of G that contains $S \cup \{v_{i-1}, v_i\}$. Set D_0 to be C and set D_{k+1} to be C' . Note that $D_i \neq D_j$ when $i < j$, because v_{i-1} is not adjacent to v_j by the minimality of k . For each $i = 0, 1, \dots, k$, the intersection of D_i and D_{i+1} contains S as well as v_i . If D_i and D_{i+1} are adjacent in $C_R(G)$ then we let P_i be the path of $C_R(G)$ consisting of D_i , D_{i+1} , and the edge between them. Otherwise D_i and D_{i+1} are not adjacent in $C_R(G)$ and the assumption on the cardinality of S means that there is a path P_i of $C_R(G)$ from D_i to D_{i+1} such that every intersection of consecutive cliques in P_i properly contains $S \cup v_i$. We concatenate the paths P_0, P_1, \dots, P_k and obtain a walk of $C_R(G)$ from C to C' . The intersection of any two consecutive cliques in this walk properly contains S . It follows that there is a path of $C_R(G)$ from C to C' with exactly the same property, and now C and C' fail to provide a counterexample after all. \square

Figure 2 illustrates Proposition 2.4. The intersection of cliques $C = \{1, 2, 3\}$ and $C' = \{3, 5, 7, 8\}$ is $\{3\} \neq \emptyset$, but C and C' are not adjacent

in $C_R(G)$. However, there is a path between C and C' in $C_R(G)$, and the intersection of any consecutive two cliques in the path properly contains $\{3\}$.




FIGURE 2. A chordal graph and its reduced clique graph.

Proposition 2.5. *Let G be a connected chordal graph. Let T be a clique tree of G . Assume that C and C' are maximal cliques of G that are adjacent in T . Then C and C' are adjacent in $C_R(G)$.*

Proof. Assume C and C' are adjacent in T , but not in $C_R(G)$. We partition the maximal cliques of G as follows. Let \mathcal{U} be the set of maximal cliques of G such that D is in \mathcal{U} if and only if the path of T from D to C does not contain C' . Similarly, define \mathcal{U}' so that D' is in \mathcal{U}' if and only if the path of T from D' to C' does not contain C . Note that every maximal clique of G is in exactly one of \mathcal{U} or \mathcal{U}' , since T is a tree. Furthermore C is in \mathcal{U} and C' is in \mathcal{U}' . Let U be the union of the cliques in \mathcal{U} , and let U' be the union of the cliques in \mathcal{U}' . Every vertex is in at least one maximal clique so $U \cup U' = V(G)$. Note that $C \subseteq U$ and $C' \subseteq U'$, so neither U nor U' is empty.

If $U \cap U' = \emptyset$, then we choose $u \in U$ and $u' \in U'$ so that u and u' are adjacent in G . (We are able to do so because G is connected.) The edge between u and u' is contained in a maximal clique. If this maximal clique is in \mathcal{U} then u' is in $U \cap U'$, and if it is in \mathcal{U}' then u is in $U \cap U'$. In either case we have a contradiction, so $U \cap U' \neq \emptyset$.

Choose an arbitrary vertex v in $U \cap U'$. Choose $D \in \mathcal{U}$ and $D' \in \mathcal{U}'$ such that v is in $D \cap D'$. Because T is a clique tree, it follows that v is contained in all the cliques belonging to the path of T from D to D' . In particular, v is contained in C and C' . Thus $U \cap U' \subseteq C \cap C'$ and $C \cap C'$ is non-empty.

Let S be $C \cap C'$. Since C and C' are not adjacent in $C_R(G)$, we can apply Proposition 2.4 and find a path P of $C_R(G)$ from C to C' , where the intersection of each pair of consecutive cliques in this path properly contains S . Since C is in \mathcal{U} and C' is in \mathcal{U}' , there is an edge of P that joins a clique $D \in \mathcal{U}$ to a clique $D' \in \mathcal{U}'$. Then $D \cap D'$ properly contains S , so we choose v in $(D \cap D') - S$. Again using the fact that T is a clique tree, we see that the

path of T from D to D' consists of cliques that contain v . In particular, v is in $C \cap C' = S$, and we have a contradiction that completes the proof. \square

It follows from Proposition 2.5 that every clique tree of G is a spanning tree of $C_R(G)$.

Proposition 2.6. *Let G be a connected chordal graph and let σ be a legitimate weighting of G . Let T be a clique tree of G . Let C and C' be maximal cliques of G that are adjacent in $C(G)$ and let P be the path of T between C and C' . The weight of any edge in P is at least $\sigma(C \cap C')$. Moreover, if C and C' are adjacent in $C_R(G)$, then at least one edge in P has weight equal to $\sigma(C \cap C')$.*

Proof. Let S be $C \cap C'$. Let P be the path of T from C to C' , and let the cliques in this path be C_0, C_1, \dots, C_n , where $C_0 = C$ and $C_n = C'$. Note that P is a path of $C_R(G)$ by Proposition 2.5. Thus any two consecutive cliques in the path have a non-empty intersection. Assume $\sigma(C_i \cap C_{i+1}) < \sigma(S)$ for some i . If S were a subset of $C_i \cap C_{i+1}$, then we would have $\sigma(S) \leq \sigma(C_i \cap C_{i+1})$ by the definition of a legitimate weighting, but this is not true. Therefore we can choose v to be a vertex in $S - (C_i \cap C_{i+1})$. Now v is a vertex of both C and C' , but the path of T between C and C' contains at least one maximal clique (either C_i or C_{i+1}) that does not contain v . This contradicts the fact that T is a clique tree. Therefore the weight of any edge in P is at least equal to $\sigma(S)$.

Now assume that C and C' are adjacent in $C_R(G)$, so that they form a separating pair. That is, there are distinct connected components of $G - S$ that contain, respectively, $C - S$ and $C' - S$. There must be maximal cliques D and D' that are adjacent in P , where $D - S$ is in the same connected component of $G - S$ as $C - S$, and $D' - S$ is not in this connected component. This means that $D \cap D'$ is contained in S . Hence $\sigma(D \cap D') \leq \sigma(S)$. The previous paragraph shows that $\sigma(D \cap D') \geq \sigma(S)$, so the result follows. \square

The proof of the next result is a straightforward adaptation of a proof given by Blair and Peyton [1, Theorem 3.6].

Lemma 2.7. *Let G be a connected chordal graph. Let σ be a legitimate weighting of G and let T be a spanning tree of $C(G)$. Then T is a clique tree of G if and only if it is a maximum-weight spanning tree of $C(G)$.*

Proof. If T is a clique tree, then for any pair of maximal cliques, C and C' , such that C and C' are adjacent in $C(G)$, the weight of the edge between C and C' is no greater than the weight of any edge in the path of T between C and C' (Proposition 2.6). It immediately follows that T has maximum weight.

For the other direction, we assume that T is a maximum-weight spanning tree. Because every chordal graph has a clique tree, and any clique tree is a spanning tree of $C_R(G)$ (and hence of $C(G)$), we can choose a clique tree T' so that T and T' have as many edges in common as possible. We can choose

an edge in T that is not in T' , because otherwise there is nothing left for us to prove. So let e be such an edge, and assume that e joins maximal cliques C and C' . There are two connected components of $T \setminus e$, one containing C and the other containing C' . Let P be the path of T' from C to C' . We let f be an edge of P which joins two cliques that are not in the same component of $T \setminus e$. Note that f is an edge of T' , and hence an edge of $C(G)$.

If $(T - e) \cup f$ is not a spanning tree of $C(G)$, then there is a path of T between the end-vertices of f that does not use e . But the end-vertices of f are in different connected components of $T \setminus e$, so $(T - e) \cup f$ is indeed a spanning tree. Similarly, if $(T' - f) \cup e$ is not a spanning tree, then there is a path of T' between C and C' that does not contain f . But P is the unique path of T' between C and C' , and f is an edge of P . So $(T - e) \cup f$ and $(T' - f) \cup e$ are both spanning trees of $C(G)$.

Applying Proposition 2.6 to the clique tree T' shows that the weight of f is at least the weight of e . Since T is a maximum-weight spanning tree, and $(T - e) \cup f$ is a spanning tree it follows that the weights on e and f must be equal. Let D and D' be the maximal cliques joined by f . Any element that is in both C and C' must be in all the cliques in P , since T' is a clique tree. This shows that $C \cap C' \subseteq D \cap D'$. If $C \cap C'$ were a proper subset of $D \cap D'$, then the definition of a legitimate weighting would mean that the weight of e is strictly less than the weight of f , which is not true. Therefore $C \cap C' = D \cap D'$.

We note that $(T' - f) \cup e$ cannot be a clique tree, since it has one more edge in common with T than T' does. Therefore we choose a vertex $v \in V(G)$ so that the maximal cliques containing v do not induce a subtree of $(T' - f) \cup e$. Let T'' be the subtree of T' induced by the maximal cliques containing v . Then f is in T'' , or else T'' would be a subtree of $(T' - f) \cup e$. This means that v is in $D \cap D' = C \cap C'$. So both C and C' are in T'' , but they are not in the same component of $T'' \setminus f$, because in that case $(T' - f) \cup e$ would contain a cycle. So e joins two vertices of T'' that are in different components of $T'' \setminus f$. Thus $(T'' - f) \cup e$ is a subtree of $(T' - f) \cup e$, and we have a contradiction that completes the proof. \square

Proof of Theorem 2.2. We have already noted that every clique tree is a spanning tree of $C_R(G)$. Let T be a clique tree of G . Then T is a maximum-weight spanning tree of $C(G)$ by Lemma 2.7. But every edge of T is an edge of $C_R(G)$, by Proposition 2.5. Since $C_R(G)$ is a subgraph of $C(G)$ it follows that T is a maximum-weight spanning tree of $C_R(G)$.

For the other direction, we let T be a maximum-weight spanning tree of $C_R(G)$. We claim that T is also a maximum-weight spanning tree of $C(G)$. To prove this claim, let e be an arbitrary edge of $C(G)$ that is not in T , let C and C' be the maximal cliques of G joined by e , and let P be the path of T that joins C and C' . If e is an edge of $C_R(G)$, then the weight of e is no greater than the weight of any edge in P , since T is a maximum-weight spanning tree of $C_R(G)$. Therefore we assume that e is

not an edge of $C_R(G)$. Now it follows from Proposition 2.4 and the definition of a legitimate weighting that there is a path from C to C' such that the edges in this path all have weight strictly greater than the weight of e . From considering, say, Kruskal's algorithm, we now see that the edges in P all have weight strictly greater than the weight of e . In either case, the weight of e does not exceed the weight of any edge in P . This implies that T is indeed a maximum-weight spanning tree of $C(G)$, and thus T is a clique tree of G by Lemma 2.7.

To complete the proof, we let e be an arbitrary edge of $C_R(G)$. We will prove that e is in a maximum-weight spanning tree of $C_R(G)$. We let C and C' be the maximal cliques joined by e . Let T be an arbitrary maximum-weight spanning tree of $C_R(G)$, so that T is a clique tree by the previous paragraph. If e is in T then we have nothing left to prove, so assume that P is the path of T joining C to C' , where P contains more than one edge. Proposition 2.6 shows that P contains an edge, f , with weight equal to the weight of e . Now $(T - f) \cup e$ is a maximum-weight spanning tree of $C_R(G)$ that contains e , and we are done. \square

From the previous arguments we can deduce further additional facts, both noted in [3]: any edge that is in $C(G)$ but not $C_R(G)$ cannot be in any maximum-weight spanning tree of $C(G)$. Secondly, $C_R(G)$ is in fact the union of all clique trees of G .

Although the next fact is incidental to our main results here, we note it for a future application in [9].

Proposition 2.8. *Let G be a connected chordal graph, and let T be a clique tree of G . Let C and C' be adjacent in T and let S be $C \cap C'$. Assume that D and D' are maximal cliques of G and the path of T from D to D' contains both C and C' . Then $D - S$ and $D' - S$ are in different connected components of $G - S$.*

Proof. Let \mathcal{U} be the family of maximal cliques of G such that D is in \mathcal{U} if and only if the path of T from D to C does not contain C' . Similarly, we let \mathcal{U}' be the family of maximal cliques where D' is in \mathcal{U}' if and only if the path of T from D' to C' does not contain C . Note that every maximal clique of G belongs to exactly one of \mathcal{U} and \mathcal{U}' . We are asserting that if $D \in \mathcal{U}$ and $D' \in \mathcal{U}'$, then $D - S$ and $D' - S$ are in different connected components of $G - S$. Assume that this fails for D and D' , where $D \cap D'$ is as large as possible. Let H be the connected component of $G - S$ that contains both $D - S$ and $D' - S$.

Let P be the path of T from D to D' . Therefore P contains both C and C' . Let v be an arbitrary vertex of $D \cap D'$. Then v is in every maximal clique that appears in P , since T is a clique tree. In particular, v is in C and C' . Thus v is in S , and this shows that $D \cap D'$ is contained in S .

Let v_0, v_1, \dots, v_k be the vertex sequence of a shortest-possible path of H from a vertex $v_0 \in D - S$ to a vertex $v_k \in D' - S$. This is an S -avoiding path, where S contains $D \cap D'$. Thus we can apply Proposition 2.3. For

$i = 1, 2, \dots, k$ we let D_i be a maximal clique of G that contains $(D \cap D') \cup \{v_{i-1}, v_i\}$. Let D_0 be D and let D_{k+1} be D' . Note that each $D_i - S$ is contained in H . This is true for D_0 and D_{k+1} by definition, and every other D_i contains the edge $v_{i-1}v_i$, which is in the path of H from v_0 to v_k . Since D_0 is in \mathcal{U} and D_{k+1} is in \mathcal{U}' , we can choose i so that D_i is in \mathcal{U} and D_{i+1} is in \mathcal{U}' . The intersection of D_i and D_{i+1} is larger than $D \cap D'$, since it contains $(D \cap D') \cup v_i$. As $D_i - S$ and $D_{i+1} - S$ are both contained in H we have a contradiction to the choice of D and D' . \square

3. THE STRUCTURE OF REDUCED CLIQUE GRAPHS

Habib and Stacho comment on the possibility of investigating the structure of graphs that are isomorphic to reduced clique graphs [6, p. 714]. In this section we make a contribution to this investigation. We start by answering an obvious question that requires a non-trivial proof.

Corollary 3.1. *Let G be a chordal graph. Then $C_R(G)$ is connected if and only if G is connected.*

Proof. Assume that H and H' are distinct connected components of G . No maximal clique of H can share a vertex with a maximal clique of H' . It follows that there be no path of $C_R(G)$ that joins two such cliques. Thus $C_R(G)$ is not connected.

The other direction is stated without proof in [6, p. 716]. Assume that G is connected. Since G is chordal it has a clique tree [4, Theorem 2], and Proposition 2.5 shows that every edge of the clique tree is an edge of $C_R(G)$. Thus $C_R(G)$ has a spanning tree, so it is connected. \square

Next we note a characterisation of clique graphs due to Szwarcfiter and Bornstein.

Theorem 3.2 ([11, Theorem 2.1]). *The graph H is isomorphic to $C(G)$ for some connected chordal graph G if and only if H has a spanning tree T such that whenever u and v are adjacent in H , the path of T from u to v induces a clique of H .*

3.1. Induced cycles. Next we observe that clique graphs can have induced cycles of any length. We will later show that this is not true for reduced clique graphs. For an integer $n \geq 3$ the *wheel graph* with n spokes is obtained from a cycle of n vertices by adding a new vertex and making it adjacent to all vertices of the cycle. Thus the wheel graph with n spokes has an induced cycle of n vertices.

Proposition 3.3. *For every integer $n \geq 3$ the wheel graph with n spokes is isomorphic to the clique graph of a chordal graph.*

Proof. This is easy to prove using Theorem 3.2, but we will give a direct construction. Start with a clique on the $n + 1$ vertices $u_0, u_1, \dots, u_{n-1}, x$. For each $i \in \mathbb{Z}/n\mathbb{Z}$, add a new vertex v_i and make it adjacent to u_i and u_{i+1} .

Call the resulting graph G . It is easy to verify that G is chordal, and its maximal cliques are $\{u_0, u_1, \dots, u_{n-1}, x\}$ along with $\{v_i, u_i, u_{i+1}\}$ for each $i \in \mathbb{Z}/n\mathbb{Z}$. The result follows. \square

Proposition 3.4. *For every even integer $n \geq 4$ the wheel graph with n spokes is isomorphic to the reduced clique graph of a chordal graph.*

Proof. The example in Figure 1 shows that this proposition is true when $n = 4$. Therefore we will assume that n is an even integer exceeding 4. Let k be $n/2$. We start with a clique on $u_0, u_1, \dots, u_{k-1}, x$. For each $i \in \mathbb{Z}/k\mathbb{Z}$, add a new vertex v_i and make it adjacent to u_i and u_{i+1} . In addition, add a new vertex z_i and make it adjacent to u_i . Let G be the resulting graph. Certainly G is chordal. The maximal cliques of G are $\{u_0, u_1, \dots, u_{k-1}, x\}$ along with $\{v_i, u_i, u_{i+1}\}$ and $\{u_i, z_i\}$ for each $i \in \mathbb{Z}/k\mathbb{Z}$. The fact that $k \geq 3$ means that $\{v_i, u_i, u_{i+1}\}$ and $\{v_j, u_j, u_{j+1}\}$ have at most one vertex in common when $i \neq j$. We can easily verify that such a pair does not form a separating pair. On the other hand, $\{u_i, z_i\}$ forms a separating pair with both $\{v_i, u_i, u_{i+1}\}$ and $\{v_{i-1}, u_{i-1}, u_i\}$. All maximal cliques form a separating pair with $\{u_0, u_1, \dots, u_{k-1}, x\}$. In this way we can check that $C_R(G)$ is isomorphic to the wheel graph with n spokes. \square

If G is the chordal graph $K_{1,4}$, then G has four maximal cliques and $C_R(G)$ is the complete graph on four vertices (which is to say, the wheel with three spokes). However, we see no way of constructing a reduced clique graph that is isomorphic to a wheel graph with a number of spokes that is odd and greater than three. In fact, we are prepared to make the following, stronger, conjecture.

Conjecture 3.5. *Let $k > 3$ be an odd integer. There is no chordal graph G such that $C_R(G)$ contains an induced cycle with exactly k vertices.*

We prove the first case of this conjecture in the following work.

Definition 3.6. Let G be a chordal graph. Let C_0, C_1, \dots, C_{n-1} be a cyclic ordering of the maximal cliques in an induced cycle of $C_R(G)$. We take the indices to be from $\mathbb{Z}/n\mathbb{Z}$, so C_i and C_j are adjacent in $C_R(G)$ if and only if $j \in \{i-1, i+1\}$. If $|C_i \cap C_{i+1}| \leq |C_j \cap C_{j+1}|$ for every $j \in \mathbb{Z}/n\mathbb{Z}$, then we say that the edge between C_i and C_{i+1} is a *minimal* edge of the cycle.

Lemma 3.7. *Let G be a chordal graph. Let C_0, C_1, \dots, C_{n-1} be a cyclic ordering of the maximal cliques in an induced cycle of $C_R(G)$, where $n \geq 4$ and the indices are from $\mathbb{Z}/n\mathbb{Z}$. Assume that the edge between C_0 and C_1 is a minimal edge of the induced cycle. Let S be $C_0 \cap C_1$ and for $i = 0, 1$ let H_i be the connected component of $G - S$ that contains $C_i - S$. Then H_0 and H_1 are distinct connected components and $C_i - S$ is contained in H_0 or H_1 for every $i \in \mathbb{Z}/n\mathbb{Z}$. Furthermore, either:*

- (i) H_0 contains all of $C_0 - S, C_2 - S, \dots, C_{n-1} - S$,
- (ii) H_1 contains all of $C_1 - S, C_2 - S, \dots, C_{n-1} - S$, or

- (iii) $n = 4$, and H_0 contains $C_0 - S$ and $C_2 - S$ while H_1 contains $C_1 - S$ and $C_3 - S$.

Proof. Note that because C_0, C_1, \dots, C_{n-1} are distinct maximal cliques of G , none of them is contained in S . Thus $C_i - S$ is non-empty for all i and is contained in H_i , a connected component of $G - S$. Because C_0 and C_1 form a separating pair, $C_0 - S$ and $C_1 - S$ are contained in different connected components of $G - S$, so H_0 and H_1 are distinct components.

Claim 3.7.1. *Assume that i and j are distinct indices in $\mathbb{Z}/n\mathbb{Z}$ such that H_i and H_j are distinct, and furthermore, C_i is adjacent in $C_R(G)$ to C_p , where $C_p - S$ is not contained in H_i and C_j is adjacent to C_q , where $C_q - S$ is not contained in H_j . Then C_i and C_j are adjacent in $C_R(G)$.*

Proof. Note that because the cycle of $C_R(G)$ is induced, p is in $\{i-1, i+1\}$ and q is in $\{j-1, j+1\}$. Note also that $C_i \cap C_p$ is contained in S . If this containment is proper then $|C_i \cap C_p| < |S| = |C_0 \cap C_1|$ and we have violated our assumption that the edge between C_0 and C_1 is minimal. Therefore C_i and C_p both contain S . The same argument shows $S \subseteq C_j \cap C_q$. Now $C_i \cap C_j$ is equal to S . Moreover $C_i - S$ and $C_j - S$ are in different components of $G - S$, so C_i and C_j form a separating pair of maximal cliques. Hence they are adjacent in $C_R(G)$. \square

We colour the cliques of C_0, C_1, \dots, C_{n-1} in the following way. For each $i \in \mathbb{Z}/n\mathbb{Z}$, if $C_i - S$ is contained in H_0 we colour C_i red, and if $C_j - S$ is in H_1 we colour C_i blue. Thus C_0 is red and C_1 is blue.

Claim 3.7.2. *Any maximal clique C_i is either red or blue.*

Proof. If the claim fails then there is some $i \in \mathbb{Z}/n\mathbb{Z} - \{0, 1\}$ such that H_i is not equal to H_0 or H_1 . We colour any clique C_j in C_0, C_1, \dots, C_{n-1} green if $C_j - S$ is contained in H_i . We know that the collections of red, blue, and green cliques are all non-empty. Therefore we can find a red clique, C_{red} , adjacent to a clique that is not red. We can similarly find C_{blue} , a blue clique that is adjacent to a non-blue clique, and C_{green} , a green clique that is adjacent to a clique that is not green. Now Claim 3.7.1 implies that C_{red} , C_{blue} , and C_{green} are adjacent to each other in $C_R(G)$. As they are three distinct vertices in an induced cycle of $C_R(G)$ with at least four vertices, this is an immediate contradiction. \square

If C_1 is the only blue clique, then statement (i) holds and we have nothing left to prove. Similarly, if C_0 is the only red clique, then (ii) holds and we are done. So we assume there are at least two red cliques and at least two blue cliques. We can choose C_{red} and C'_{red} to be distinct red cliques that are adjacent to blue cliques, and we can choose C_{blue} and C'_{blue} to be two distinct blue cliques that are adjacent to red cliques. Now Claim 3.7.1 implies that C_{red} and C'_{red} are adjacent to both C_{blue} and C'_{blue} . Thus the four cliques induce a cycle in $C_R(G)$. This is impossible if $n \geq 5$, so we conclude that

$n = 4$. Now C_0 is a red clique and it is adjacent to two blue cliques. Thus C_1 and C_3 are blue, C_2 is red, and we are finished. \square

We next establish the first case of Conjecture 3.5.

Lemma 3.8. *There is no chordal graph G such that $C_R(G)$ has an induced cycle with exactly five vertices.*

Proof. Assume otherwise and let G be a chordal graph such that $C_R(G)$ contains an induced cycle with five vertices. Let C_0, C_1, C_2, C_3, C_4 be the maximal cliques in this cycle, where the indices are from $\mathbb{Z}/5\mathbb{Z}$ and C_i is adjacent to C_j if and only if $j \in \{i-1, i+1\}$. By adding a constant to these indices as necessary, we may assume that

$$|C_0 \cap C_1| \leq |C_i \cap C_{i+1}|$$

for all $i \in \mathbb{Z}/5\mathbb{Z}$, so that the edge between C_0 and C_1 is a minimal edge of the cycle. Let S be $C_0 \cap C_1$. Note that S is non-empty.

Now we apply Lemma 3.7. By applying the permutation $\rho: i \mapsto 1-i$ as necessary, we may assume that statement (ii) in Lemma 3.7 applies. Therefore we let H_0 and H_1 be connected components of $G - S$ such that H_0 contains $C_0 - S$ and H_1 contains $C_1 - S, C_2 - S, C_3 - S,$ and $C_4 - S$.

Claim 3.8.1. $C_0 \cap C_4 = S = C_0 \cap C_1$.

Proof. Because $C_0 - S$ and $C_4 - S$ are contained in different components of $G - S$, it follows that $C_0 \cap C_4 \subseteq S$. All we have left to prove is that this containment is not proper. If it were proper, then we would contradict the assumption that the edge between C_0 and C_1 is minimal. \square

Claim 3.8.2. *Neither C_2 nor C_3 contains S .*

Proof. Note that $C_0 \cap C_2 \subseteq S$ because $C_0 - S$ and $C_2 - S$ are contained in different components of $G - S$. Certainly any path from a vertex of $C_0 - C_2$ to a vertex of $C_2 - C_0$ must use a vertex of S . If $C_0 \cap C_2 = S$, then C_0 and C_2 form a separating pair, so C_0 and C_2 are adjacent in $C_R(G)$. This contradicts the fact that C_0 and C_2 are non-consecutive vertices in an induced cycle. The same argument shows that C_3 does not contain S . \square

Claim 3.8.3. $C_2 \cap C_4 \subseteq C_1$ and $C_3 \cap C_1 \subseteq C_4$.

Proof. Assume that x is a vertex of $C_2 \cap C_4$ that is not in C_1 . By Claim 3.8.2 we can let y be a vertex in $S - C_2$. Thus y is in $C_1 - C_2$. So x is in $C_2 - C_1$ and y is in $C_1 - C_2$. Claim 3.8.1 implies that y is in C_4 . As x is also in C_4 we see that x and y are adjacent. Because C_1 and C_2 are adjacent in $C_R(G)$ they have a non-empty intersection, but now the edge xy shows that C_1 and C_2 do not form a separating pair and we have a contradiction. A symmetric argument shows $C_3 \cap C_1 \subseteq C_4$. \square

Claim 3.8.4. C_2 contains a vertex of $C_1 - C_4$ and C_3 contains a vertex of $C_4 - C_1$.

Proof. By symmetry it suffices to prove the first statement. Assume that C_2 contains no vertex of $C_1 - C_4$. Because C_1 and C_2 are adjacent in $C_R(G)$, they have at least one vertex in common. By our assumption, no vertex of $C_1 \cap C_2$ is in $C_1 - C_4$, so any such vertex must be in $C_1 \cap C_4$. Therefore C_2 and C_4 are not disjoint. Since C_2 and C_4 are not adjacent in $C_R(G)$, we can let P be a $(C_2 \cap C_4)$ -avoiding path from a vertex $x \in C_2 - C_4$ to $y \in C_4 - C_2$.

Our assumption means that x is not in C_1 , so it is in $C_2 - C_1$. Our assumption and Claim 3.8.3 imply that $C_2 \cap C_4 = C_2 \cap C_1$. Therefore P is a $(C_2 \cap C_1)$ -avoiding path. But Claim 3.8.2 shows that we can choose a vertex z in $S - C_2$. Thus z is in $C_1 - C_2$ and Claim 3.8.1 shows that z is in C_4 . Assuming that z and y are not equal, they are adjacent, as both are in C_4 . By appending (if necessary) the edge yz to the end of P we obtain a $(C_1 \cap C_2)$ -avoiding path from a vertex in $C_2 - C_1$ to a vertex in $C_1 - C_2$. Hence C_1 and C_2 do not form a separating pair and this contradicts the fact that they are adjacent in $C_R(G)$. \square

Claim 3.8.5. *Either $C_2 \cap (C_1 \cap C_4) \subseteq C_3$ or $C_3 \cap (C_1 \cap C_4) \subseteq C_2$.*

Proof. Note that $C_2 \cap C_3$ is non-empty, since C_2 and C_3 are adjacent in $C_R(G)$. If the claim fails, then we choose $x \in (C_2 \cap C_1 \cap C_4) - C_3$ and $y \in (C_3 \cap C_1 \cap C_4) - C_2$. Now x and y are both in $C_1 \cap C_4$, so they are adjacent. Moreover x is in $C_2 - C_3$ and y is in $C_3 - C_2$. Thus C_2 and C_3 do not form a separating pair and we have a contradiction. \square

By using Claim 3.8.5, we will assume that $C_2 \cap (C_1 \cap C_4)$ is a subset of C_3 . The other outcome from Claim 3.8.5 yields to a symmetric argument. Using Claim 3.8.2 we choose a vertex $x \in S$ that is not in C_3 . Note that Claim 3.8.1 implies that S is contained in $C_1 \cap C_4$. So x is in $(C_1 \cap C_4) - C_3$. The assumption $C_2 \cap (C_1 \cap C_4) \subseteq C_3$ implies that x is not in C_2 .

By Claim 3.8.4 we can also choose y in $C_2 \cap (C_1 - C_4)$ and z in $C_3 \cap (C_4 - C_1)$. Because y is in $C_1 - C_4$ and Claim 3.8.3 says that $C_3 \cap C_1$ is a subset of C_4 , it follows that y is not in C_3 . Therefore y belongs to $C_2 - C_3$. A symmetric argument shows that z is in $C_3 - C_2$. Now x and y are adjacent as they are both in C_1 , and x and z are adjacent as they are both in C_4 . Note that $C_2 \cap C_3$ is non-empty as C_2 and C_3 are adjacent in $C_R(G)$. But the path with vertex sequence y, x, z is $(C_2 \cap C_3)$ -avoiding, so C_2 and C_3 do not form a separating pair. This final contradiction completes the proof. \square

Lemma 3.8 shows that the class of reduced clique graphs is contained in the class of graphs with no length-five induced cycle. We next show that this containment is proper.

Proposition 3.9. *Let $n \geq 4$ be an integer. There is no chordal graph G such that either $C(G)$ or $C_R(G)$ is a cycle with n vertices.*

Proof. Let H be a cycle with at least four vertices. Any spanning tree of H is a Hamiltonian path. The end vertices of this path are adjacent in H , but the path of the spanning tree between these vertices does not induce a

clique in H . Therefore H is not isomorphic to $C(G)$ for any chordal graph G by Theorem 3.2.

We turn to reduced chordal graphs. Assume for a contradiction that G is a chordal graph with C_0, C_1, \dots, C_{n-1} as its list of maximal cliques, where the indices are from $\mathbb{Z}/n\mathbb{Z}$ for some $n \geq 4$, and C_i is adjacent to C_j in $C_R(G)$ if and only if $j \in \{i-1, i+1\}$. We can assume without loss of generality that the edge between C_0 and C_1 is a minimal edge of $C_R(G)$. Let S be $C_0 \cap C_1$. Assume that statement (iii) in Lemma 3.7 holds. Thus $n = 4$ and there are distinct connected components, H_0 and H_1 , of $G - S$ such that H_0 contains $C_0 - S$ and $C_2 - S$ while H_1 contains $C_1 - S$ and $C_3 - S$. Note that $C_0 \cap C_3 \subseteq S$, and in fact $C_0 \cap C_3$ is equal to S , or else the minimality of the C_0 - C_1 edge is contradicted.

Either $C_0 \cap C_2$ is empty, or it is not. In the latter case, we can apply Proposition 2.4 to C_0 and C_2 . We see that either $C_0 \cap C_1$ or $C_0 \cap C_3$ properly contains $C_0 \cap C_2$. By symmetry, we can assume $C_0 \cap C_2$ is a proper subset of $C_0 \cap C_1 = S$. Thus $C_0 - S$ and $C_2 - S$ are disjoint sets. We can let P be a shortest-possible path of H_0 from a vertex of $C_0 - S$ to a vertex of $C_2 - S$. On the other hand, if $C_0 \cap C_2$ is empty, then $C_0 - S$ and $C_2 - S$ are again disjoint subsets in H_0 , so we again let P be a shortest-possible path of H_0 from $C_0 - S$ to $C_2 - S$. In either case, P contains exactly one vertex of C_0 and exactly one vertex of C_2 . Then P must contain at least one edge, and this edge is in a maximal clique that is equal to neither C_0 nor C_2 . Nor can this maximal clique be C_1 or C_3 , because any edge of P is contained in H_0 . So we have a contradiction in the case that (iii) in Lemma 3.7 holds.

Now we assume that either (i) or (ii) holds. By applying the permutation $\rho: i \mapsto 1 - i$ as necessary, we will assume that H_0 and H_1 are distinct connected components of $G - S$, and that H_0 contains $C_0 - S$ while H_1 contains $C_i - S$ for $i = \{1, 2, \dots, n-1\}$. By the same argument as earlier, we can see that C_{n-1} contains S , or else the choice of the $C_0 - C_1$ edge is contradicted.

Now $C_1 \cap C_{n-1}$ contains S , and C_1 and C_{n-1} are non-adjacent in $C_R(G)$. We apply Proposition 2.4 and see that there is a path of $C_R(G)$ from C_1 to C_{n-1} such that every intersection of consecutive cliques in the path properly contains $C_1 \cap C_{n-1}$. This path is either C_1, C_0, C_{n-1} , or it is C_1, C_2, \dots, C_{n-1} . Assume the former. Then $C_1 \cap C_0 = S$ properly contains $C_1 \cap C_{n-1} \supseteq S$ and we have a contradiction. Hence any intersection of consecutive cliques in C_1, C_2, \dots, C_{n-1} properly contains $C_1 \cap C_{n-1}$, and hence contains S . It follows that C_2 contains S and thus $C_0 \cap C_2$ is non-empty.

Since $C_0 - S$ and $C_2 - S$ are contained in different components of $G - S$, any path of a vertex from $C_0 - C_2$ to a vertex of $C_2 - C_0$ must contain a vertex of $S = C_0 \cap C_2$. Thus C_0 and C_2 form a separating pair in G , and hence they are adjacent in $C_R(G)$, which is a contradiction. \square

3.2. Clique graphs vs. reduced clique graphs. Consider the classes $\{C(G)\}$ and $\{C_R(G)\}$, where G ranges over all chordal graphs. Proposition 3.3 and Lemma 3.8 show that the wheel with five spokes is isomorphic to a graph in the former class but not the latter. Is there a graph that is isomorphic to a graph in the latter class but not the former? We will show that the answer is, once again, yes. Recall that if G and G' are disjoint graphs, then $G \boxtimes G'$ is obtained from the union of G and G' by making every vertex of G adjacent to every vertex of G' . We use P_n to denote the path with n vertices.

Lemma 3.10. *Let $m, n \geq 1$ be integers. Then $P_m \boxtimes P_n$ is isomorphic to the reduced clique graph of a chordal graph. If $n \geq 22$, then $P_n \boxtimes P_n$ is not isomorphic to the clique graph of a chordal graph.*

Proof. Let G be the graph obtained from the disjoint union of P_{m+1} and P_{n+1} by adding a new vertex that is adjacent to every vertex of the disjoint union. It is easy to confirm that G is chordal, and that $C_R(G)$ is isomorphic to $P_m \boxtimes P_n$.

For the second statement, we let H be a graph with disjoint induced paths $P_u = u_0, u_1, \dots, u_{n-1}$ and $P_v = v_0, v_1, \dots, v_{n-1}$, where $n \geq 22$ and every u_i is adjacent to every v_j . Thus H is isomorphic to $P_n \boxtimes P_n$. We will assume for a contradiction that H is isomorphic to $C(G)$ for some chordal graph G . Because $C(G)$ is connected it follows easily that G is connected, so we can apply Theorem 3.2 and deduce that H has a spanning tree T , where the path of T from u to v induces a clique of H whenever u and v are adjacent in H .

Claim 3.10.1. *Let i and j be integers satisfying $0 < i, j < n-1$. The path of T from u_i to v_j is contained in one of: $\{u_i, u_{i+1}, v_j, v_{j+1}\}$, $\{u_i, u_{i+1}, v_{j-1}, v_j\}$, $\{u_{i-1}, u_i, v_j, v_{j+1}\}$, $\{u_{i-1}, u_i, v_{j-1}, v_j\}$.*

Proof. Let P be the path of T from u_i to v_j . Since u_i is adjacent to v_j it follows that P induces a clique of H . As u_i is not adjacent to any of the vertices in $u_0, \dots, u_{i-2}, u_{i+2}, \dots, u_{n-1}$, it follows that the vertices of P that are in P_u belong to $\{u_{i-1}, u_i, u_{i+1}\}$. Similarly, the vertices of P that are in P_v belong to $\{v_{j-1}, v_j, v_{j+1}\}$. But u_{i-1} is not adjacent to u_{i+1} , so P does not contain both. The claim follows by symmetry. \square

Claim 3.10.1 implies that the path of T between u_i and v_j has at most three edges.

Let P be a longest-possible path of T and let p_0, p_1, \dots, p_{k-1} be the vertices of P . For $i = 0, 1, \dots, k-1$, let U_i be the set of vertices in P_u such that u is in U_i if and only if the shortest path of T from u to a vertex in P contains p_i . We define V_i to be the analogous set of vertices in P_v . Note that $(U_0, U_1, \dots, U_{k-1})$ is a partition of the vertices of P_u , and $(V_0, V_1, \dots, V_{k-1})$ is a partition of the vertices of P_v .

Claim 3.10.2. *Either*

$$\max\{|U_i|: 0 \leq i \leq k-1\} \leq 3 \quad \text{or} \quad \max\{|V_i|: 0 \leq i \leq k-1\} \leq 3.$$

Proof. Assume for a contradiction that $|U_i| \geq 4$ and $|V_j| \geq 4$. Let p and q , respectively, be the smallest (largest) integers such that $u_p, u_q \in U_i$. Then $q-1 > p+1$ because $|U_i| \geq 4$. In the same way, let s and t be the smallest (largest) integers such that $v_s, v_t \in V_j$. Then $t-1 > s+1$. It is simple to see from Claim 3.10.1 that the path of T from u_p to v_s has no vertex in common with the path of T from u_q to v_t . But this contradicts the fact that both paths contain p_i and p_j . \square

By using Claim 3.10.2, we will assume without loss of generality that $|V_i| \leq 3$ for each $i = 0, 1, \dots, k-1$. Since $(U_0, U_1, \dots, U_{k-1})$ is a partition of the vertices in P_u we can choose i so that U_i contains a vertex x . We claim that if $j \leq i-4$ or $j \geq i+4$, then $V_j = \emptyset$. If this fails, then the path of T from a vertex in V_j to x contains at least four edges of P . But this contradicts our earlier conclusion that any path of T from a vertex of P_u to a vertex of P_v contains at most three edges. So now the vertices of P_v belong to

$$V_{i-3} \cup V_{i-2} \cup \dots \cup V_{i+2} \cup V_{i+3}$$

and this union has cardinality at most 7×3 . Thus P_v contains at most 21 vertices and this contradicts $n \geq 22$. \square

4. CONCLUSIONS AND OPEN PROBLEMS

We restate Conjecture 3.5 here.

Conjecture 4.1. *Let $k > 3$ be an odd integer. There is no chordal graph G such that $C_R(G)$ contains an induced cycle with exactly k vertices.*

So far as we have been able to tell, every chordal graph is isomorphic to both a clique graph, and to a reduced clique graph. We conjecture this holds generally.

Conjecture 4.2. *Let H be a chordal graph. There are chordal graphs G and G' such that H is isomorphic to both $C(G)$ and $C_R(G')$.*

Szwarcfiter and Bornstein present a polynomial-time algorithm for deciding whether a given graph is isomorphic to $C(G)$ for some chordal graph G [11]. Their techniques do not obviously extend to recognising reduced clique graphs. Nonetheless, we will make the following conjecture.

Conjecture 4.3. *There is a polynomial-time algorithm for deciding whether a given graph is isomorphic to $C_R(G)$ for some chordal graph G .*

More informally, we ask if there is a structural description for reduced clique graphs that is analogous to Theorem 3.2.

5. ACKNOWLEDGMENTS

We thank the anonymous referee for their careful work, and for providing the construction in Proposition 3.4.

REFERENCES

- [1] Jean R. S. Blair and Barry Peyton, *An introduction to chordal graphs and clique trees*, Graph theory and sparse matrix computation, IMA Vol. Math. Appl., vol. 56, Springer, New York, 1993, pp. 1–29.
- [2] Peter Buneman, *A characterisation of rigid circuit graphs*, Discrete Math. **9** (1974), 205–212.
- [3] Philippe Galinier, Michel Habib, and Christophe Paul, *Chordal graphs and their clique graphs*, Graph-theoretic concepts in computer science (Aachen, 1995), Lecture Notes in Comput. Sci., vol. 1017, Springer, Berlin, 1995, pp. 358–371.
- [4] Fănică Gavril, *The intersection graphs of subtrees in trees are exactly the chordal graphs*, J. Combinatorial Theory Ser. B **16** (1974), 47–56.
- [5] Michel Habib and Vincent Limouzy, *On some simplicial elimination schemes for chordal graphs*, DIMAP Workshop on Algorithmic Graph Theory, Electron. Notes Discrete Math., vol. 32, Elsevier Sci. B. V., Amsterdam, 2009, pp. 125–132.
- [6] Michel Habib and Juraj Stacho, *Reduced clique graphs of chordal graphs*, European J. Combin. **33** (2012), no. 5, 712–735.
- [7] Terry A. McKee, *Minimal weak separators of chordal graphs*, Ars Combin. **101** (2011), 321–331.
- [8] Yasuko Matsui, Ryuhei Uehara, and Takeaki Uno, *Enumeration of the perfect sequences of a chordal graph*, Theoret. Comput. Sci. **411** (2010), no. 40–42, 3635–3641.
- [9] Dillon Mayhew and Andrew Probert, *Supersolvable saturated matroids and chordal graphs*, Adv. in Appl. Math. **153** (2024), Paper No. 102616, 36.
- [10] Donald J. Rose, *Triangulated graphs and the elimination process*, J. Math. Anal. Appl. **32** (1970), 597–609.
- [11] Jayme L. Szwarcfiter and Claudson F. Bornstein, *Clique graphs of chordal and path graphs*, SIAM J. Discrete Math. **7** (1994), no. 2, 331–336.

6. STATEMENTS & DECLARATIONS

The first author was supported by a Rutherford Discovery Fellowship, managed by Royal Society Te Apārangi.

The authors have no relevant financial or non-financial interests to disclose.

All authors contributed to the study conception and design. Much of the material appeared previously in the PhD thesis of the second author. All authors read and approved the final manuscript.

7. DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

SCHOOL OF COMPUTING, UNIVERSITY OF LEEDS, UNITED KINGDOM

Email address: `d.mayhew@leeds.ac.uk`

Email address: `andrew.mprobert@gmail.com`